

Chapter 13

Complex numbers

$$i \triangleq \sqrt{-1}$$

$$e^{i\theta} = \cos(\theta) + i \sin(\theta)$$

Summary: Tools and use for complex numbers (see examples in Hw 8, 9)

This chapter provides algebraic tools for complex numbers (+, *, /, $\sqrt{\quad}$, and powers).¹
Complex numbers are useful in many aspects of dynamic systems, including:

Solving 2 nd , 3 rd , and higher-order ODEs	Root locus	
Control system design	Frequency response	Eigen-analysis

Motivating questions

- The following proof involves the imaginary number i defined as $i \triangleq \sqrt{-1}$ and shows the surprising conclusion that $1 = -1$. Circle the incorrect step in the proof and explain your reasoning.

$$1 = \sqrt{1} = \sqrt{(-1)^2} = \sqrt{-1} * \sqrt{-1} = i * i = i^2 = -1$$

When x is negative, $(x^2)^{(1/2)} \neq (x^{1/2})^2$. In general, $(x^a)^b \neq (x^b)^a$.

- Find all real and/or complex numbers that can appear on the right-hand side of the equal signs.²

$$1^4 = (1 e^{2n\pi i})^4_{n=0,1,2,\dots} = e^{8n\pi i} = \cos(8n\pi) + i \sin(8n\pi) = 1$$

$$1^{1/4} = (1 e^{2n\pi i})^{1/4}_{n=0,1,2,\dots} = e^{\frac{n\pi}{2} i} = \cos\left(\frac{n\pi}{2}\right) + i \sin\left(\frac{n\pi}{2}\right) = \pm 1, \pm i$$

$$1^{1/3} = (1 e^{2n\pi i})^{1/3}_{n=0,1,2,\dots} = e^{\frac{2n\pi}{3} i} = \cos\left(\frac{2n\pi}{3}\right) + i \sin\left(\frac{2n\pi}{3}\right) = 1, -0.5 \pm 0.866 i$$

¹Complex numbers are said to be “**closed**” under addition, subtraction, negation, multiplication, division, and exponentiation because when these operations are performed on complex numbers, only complex numbers result. Complex numbers are said to be “**algebraically closed**” because polynomial equations with complex number coefficients can only produce complex numbers. Real numbers are not closed under exponentiation. For example, $-4^{0.5} = \sqrt{-4} = \pm 2i$ produces a complex (not real) number. Real numbers are not algebraically closed. For example, although the polynomial equation $x^2 + 2x + 5 = 0$ has real coefficients, its roots are the complex numbers $x = -1 \pm 2i$.

²Answers at www.MotionGenesis.com \Rightarrow [Textbooks](#) \Rightarrow [Resources](#). Alternate: $1^{1/4} = \sqrt[4]{1} = \sqrt{\pm 1} = \pm 1$ or $\pm i$.